

On the Root Ambiguity in the Complete Solution to the Most General Fifth Degree Polynomial

Richard Drociuk

February 1, 2008

1 Abstract

Starting from the solution to Bring's equation the root ambiguity is removed from the solution to the quintic equation. This gives the five complex roots of the quintic equation as indicated by Gauss's Fundamental Theorem of Algebra.

2 Introduction

In the previous paper[Drociuk,1], the solution to the quintic was given, but the root ambiguity was not correctly removed. This problem arises because Ferrari's method for solving the quartic equation does not introduce all possible roots. Instead if one uses a Tshirnhaussen transformation as demonstrated[Drociuk,1], to the quartic equation, the five correct roots of the quintic equation,

$$x^5 + mx^4 + nx^3 + px^2 + qx + r = 0 \quad (1)$$

are selected for arbitrary coefficients m , n , p , q , and r . The roots are selected from all possible roots using conditional loops to an arbitrary precision, ϵ .

3 Ambiguity in the Quartic Solution

The quartic equation,

$$x^4 + a_3x^3 + a_2x^2 + a_1x + a_0 = 0 \quad (2)$$

is transformed with,

$$Tsh2 = x^3 + b_2x^2 + b_1x + b_0 + y_n \quad (3)$$

to the quadratic in y_n^2 ,

$$y_n^4 + B_2y_n^2 + B_0 = 0 \quad (4)$$

whose four roots are,

$$y_1 = \frac{1}{2}(-2B_2 + 2(B_2^2 - 4B_0)^{\frac{1}{2}})^{\frac{1}{2}} \quad (5)$$

$$y_2 = -\frac{1}{2}(-2B_2 + 2(B_2^2 - 4B_0)^{\frac{1}{2}})^{\frac{1}{2}} \quad (6)$$

$$y_3 = \frac{1}{2}(-2B_2 - 2(B_2^2 - 4B_0)^{\frac{1}{2}})^{\frac{1}{2}} \quad (7)$$

$$y_4 = -\frac{1}{2}(-2B_2 - 2(B_2^2 - 4B_0)^{\frac{1}{2}})^{\frac{1}{2}} \quad (8)$$

where,

$$\begin{aligned} B_0 = & 3b_0^2a_3^2a_0 + b_0^3b_2a_3^2 + b_0^2b_2^2a_2^2 - 2b_0^3b_2a_2 + 3a_3a_2b_0^3 + b_0b_2^3a_1^2 - b_0^3b_1a_3 + 4b_0b_2a_0^2 + \\ & 4b_0^2b_1a_0 + b_0^2b_1^2a_2 - 2b_0^2b_1a_2^2 - b_2^3a_0^2a_3 - 2a_0b_1^3a_2 + a_3^2a_0b_1^3 - b_2a_0^2a_1 - 2b_1a_0^2a_2 + b_1a_0a_1^2 + \\ & a_0b_1^2a_2^2 + a_1b_0^2b_2a_2 - 3b_0a_3a_0^2 + 2a_1b_0^2b_2a_3^2 - 3a_1a_3a_2b_0^2 + b_0^2a_3^2a_2b_1 - 5b_0^2b_2a_3a_0 - \\ & b_0^2a_3a_2^2b_2 + 3b_0^2b_1b_2a_1 - 2b_0^2b_2^2a_3a_1 - 2b_0b_2^3a_0a_2 - 4b_0b_2a_0b_1^2 - 2b_0b_2a_0a_2^2 + 2b_0a_1b_1^2a_2 + \\ & b_0b_2^2a_0a_1 - b_0a_3^2a_1b_1^2 + 2b_0a_3a_1^2b_1 - b_0b_2^2a_3a_1^2 + b_0b_2a_2a_1^2 - b_0a_1b_1a_2^2 + 3b_0a_1a_2a_0 - \\ & 3b_0b_1b_2a_1^2 - 5b_0a_1b_1a_0 + 3b_1b_2a_3a_0^2 + a_0b_0b_1^2a_3 + a_0b_1a_3a_2b_0 - a_0b_1b_2a_2a_1 - a_0b_2a_2b_1^2a_3 + \\ & 2b_1^2a_0^2 + b_4^2a_0^2 + b_1^4a_0 - b_0b_1^3a_1 + a_0^2b_2^2a_2 - 4b_2^2a_0^2b_1 - 3a_1b_0^3 + 3b_0^2a_1^2 - b_0a_1^3 - \\ & 3b_0^2a_2a_0 + b_0^4 + 2b_2^2a_0b_0^2 - b_0^3a_3^2 + b_0^2a_3^2 - a_1b_0^2b_1a_3 + a_0^3 - b_0^2b_2a_2b_1a_3 - b_0a_1b_2a_3a_0 + \\ & b_0b_2a_1b_1^2a_3 + 2b_0a_0b_2^2a_3a_2 - 3b_0a_0b_1b_2a_3^2 + 4b_0b_2a_0b_1a_2 + 3b_0b_2^2a_0b_1a_3 - b_0b_1a_1b_2^2a_2 + \\ & b_0a_1b_2a_2b_1a_3 + 3b_1^2a_0b_2a_1 - 2b_1^2a_0a_1a_3 - b_1a_0b_2^2a_1 + b_1^2a_0b_2^2a_2 + b_2^2a_0b_1a_3a_1 - \\ & b_2a_0b_1^3a_3 \end{aligned} \quad (9)$$

and

$$\begin{aligned} B_2 = & a_3^2a_2b_1 + 2a_1b_2a_3^2 - 6b_0b_2a_2 - 5b_2a_3a_0 + 4b_1a_0 - 3a_2a_0 + 3a_3^2a_0 - 3a_3a_2a_1 - \\ & 3b_0a_3^3 + 6b_0^2 - 9b_0a_1 + b_2^2a_2^2 + 3a_1^2 - 2b_1a_2^2 + b_1^2a_2 - b_2a_2b_1a_3 - a_3a_2^2b_2 + a_2^3 + \\ & 3b_1b_2a_1 - a_1b_1a_3 - 3b_0b_1a_3 + 3b_0b_2a_3^2 + 2a_0b_2^2 + 9a_3a_2b_0 + b_2a_2a_1 - 2b_2^2a_3a_1 \end{aligned} \quad (10)$$

The coefficients of the Tshirnhaussen transformation (3) are,

$$\begin{aligned} b_0 = & \frac{(36b_{01}b_{02}b_{03} - 108b_{00}b_{03}^2 - 8b_{02}^3 + 12(3)^{\frac{1}{2}}(4b_{01}^3b_{03} - b_{01}^2b_{02}^2 - 18b_{01}b_{02}b_{03}b_{00} + 27b_{00}^2b_{03}^2 + 4b_{00}b_{02}^3)^{\frac{1}{2}}b_{03})^{\frac{1}{3}}}{6b_{03}} \\ & - \frac{2}{3} \frac{(3b_{01}b_{03} - b_{02}^2)}{(b_{03}(36b_{01}b_{02}b_{03} - 108b_{00}b_{03}^2 - 8b_{02}^3 + 12(3)^{\frac{1}{2}}(4b_{01}^3b_{03} - b_{01}^2b_{02}^2 - 18b_{01}b_{02}b_{03}b_{00} + 27b_{00}^2b_{03}^2 + 4b_{00}b_{02}^3)^{\frac{1}{2}}b_{03})^{\frac{1}{3}} - \frac{1}{3}\frac{b_{02}}{b_{03}})} \end{aligned} \quad (11)$$

choose,

$$b_1 = 0 \quad (12)$$

and

$$b_2 = \frac{(a_3^3 + 3a_1 - 4b_0 + b_1a_3 - 3a_3a_2)}{(a_3^2 - 2a_2)} \quad (13)$$

the coefficients b_{00} , b_{01} , b_{02} and b_{03} are given by

$$b_{00} = (a_0^2 a_3^7 + 20a_2^3 a_1^3 + 2a_3^6 a_1^3 + 18a_3^3 a_1^4 - 36a_1 a_2^3 a_0 a_3^2 + 150a_3 a_1^2 a_2^2 a_0 + 29a_1 a_2^2 a_3^4 a_0 - 54a_3^3 a_1^2 a_0 a_2 - 4a_3^6 a_0 a_1 a_2 - 48a_3^2 a_0^2 a_1 a_2 + 27a_1^5 - a_3^5 a_2^2 a_1^2 + 28a_3^3 a_0^2 a_2^2 - 24a_2^3 a_0^2 a_3 + 48a_2^2 a_0^2 a_1 + 24a_2^5 a_0 a_3 + 12a_3^4 a_0^2 a_1 - 10a_3^5 a_0^2 a_2 - 14a_3^3 a_0 a_2^4 + 2a_3^5 a_0 a_2^3 - 12a_3 a_2^4 a_1^2 + 7a_3^3 a_2^3 a_1^2 - 48a_1 a_2^4 a_0 + 3a_3^5 a_1^2 a_0 + 21a_2^2 a_1^3 a_3^2 - 15a_2 a_1^3 a_3^4 - 72a_1^3 a_2 a_0 + 9a_1^3 a_3^2 a_0 - 63a_3 a_1^4 a_2)/(a_3^2 - 2a_2)^3 \quad (14)$$

$$b_{01} = (-10a_3^6 a_1^2 - 84a_3^3 a_1^3 + 80a_0 a_2^4 - 76a_2^3 a_1^2 - 64a_0^2 a_2^2 - 2a_3^4 a_2^4 + 12a_2^3 a_5 - 16a_3^4 a_0^2 - 108a_1^4 - 16a_2^6 + 120a_3^3 a_1 a_2 a_0 - 344a_3 a_1 a_2^2 a_0 + 16a_2^3 a_3^2 a_0 - 10a_3^5 a_1 a_0 - 24a_2^2 a_3^4 a_0 - 74a_2^2 a_1^2 a_3^2 + 70a_2 a_1^2 a_3^4 + 72a_3 a_2^4 a_1 - 52a_3^3 a_2^3 a_1 + 8a_3^5 a_2^2 a_1 + 4a_3^6 a_0 a_2 + 64a_2^3 a_0^2 a_2 240a_3 a_1^3 a_2 + 12a_2^2 a_0 a_1^2 + 192a_0 a_2 a_1^2)/(a_3^2 - 2a_2)^3 \quad (15)$$

$$b_{02} = (-64a_3^3 a_2 a_0 + 8a_3^5 a_0 + 16a_1 a_3^6 - 104a_1 a_3^4 a_2 + 112a_2^2 a_1 a_2^2 - 64a_3 a_2^4 + 128a_1^2 a_3^3 - 8a_3^5 a_2^2 - 304a_3 a_2 a_1^2 + 144a_1^3 + 64a_1 a_2^3 + 48a_3^3 a_2^2 + 192a_0 a_3 a_2^2 - 80a_1 a_2^2 a_0 - 128a_1 a_2 a_0)/(a_3^2 - 2a_2)^3 \quad (16)$$

$$b_{03} = (-64a_3^2 a_2^2 + 48a_3^4 a_2 - 8a_3^6 + 64a_2^2 a_0 - 64a_1^2 + 128a_3 a_2 a_1 - 64a_1 a_3^3)/(a_3^2 - 2a_2)^3 \quad (17)$$

Now the cubic(3) is solved for its three roots,

$$x_{1n} = \frac{\Delta_n - (2b_1 - \frac{2}{3}b_2^2)}{6\Delta_n - \frac{1}{3}b_2} \quad (18)$$

$$x_{2n} = -\frac{1}{12}\Delta_n + \frac{b_1 - \frac{1}{3}b_2^2}{\Delta_n} - \frac{1}{3}b_2 + i\sqrt{3}(\frac{1}{12}\Delta_n + \frac{b_1 - \frac{1}{3}b_2^2}{\Delta_n}) \quad (19)$$

$$x_{3n} = -\frac{1}{12}\Delta_n + \frac{b_1 - \frac{1}{3}b_2^2}{\Delta_n} - \frac{1}{3}b_2 - i\sqrt{3}(\frac{1}{12}\Delta_n + \frac{b_1 - \frac{1}{3}b_2^2}{\Delta_n}) \quad (20)$$

$$\Delta_n = (36b_1 b_2 - 108y_1 - 108b_0 - 8b_2^3 + 12(12b_1^3 - 3b_1^2 b_2^2 - 54b_1 b_2 y_n - 54b_1 b_2 b_0 + 81y_n^2 + 162y_n b_0 + 12y_n b_2^3 + 81b_0^2 + 12b_0 b_2^3)^{\frac{1}{2}})^{\frac{1}{3}} \quad (21)$$

By substituting the values of y_n , from equations (5) to (8), into equations (18) to (20), one can generate twelve possible roots, x_{mn} with $m \in (1, 2, 3)$ and $n \in (1, 2, 3, 4)$. Only four roots satisfy equation (2), they are found by direct substitution. So for all complex coefficients a_0 , a_1 , a_2 and a_3 in equation (2), the four roots of the quartic are contained in x_{mn} .

4 Ambiguity in the Quintic Solution

Recall in [Drociuk,1], the coefficients of the quartic Tshirnhausen Transformation,

$$Tsh1 = x^4 + dx^3 + cx^2 + bx + a + y \quad (22)$$

and let the coefficients of equation (2) be,

$$a_0 = a + y \quad (23)$$

$$a_1 = b \quad (24)$$

$$a_2 = c \quad (25)$$

$$a_3 = d \quad (26)$$

and calculate the twelve x_{mn} using the equations of the previous section. Then using nested “for loops” and “if-then” statements on Maple, x_{mn} is substituted into both the quintic (1) and the quartic (22). When both these equations are made less than ϵ , we have obtained the first root of the quintic equation (1), let it be r_1 . ϵ can be made arbitrarily close to zero. Let $\epsilon = 10^{-100}$, you may chose it to be zero if you want to wait or have a faster computer. Now r_1 is then factored out of the quintic (1), leaving a quartic equation to be solved with the following coefficients,

$$a_0 = q + r_1p + r_1^2n + mr_3 + r_1^4 \quad (27)$$

$$a_1 = p + r_1n + r_1^2m + r_1^3 \quad (28)$$

$$a_2 = n + mr_1 + r_1^2 \quad (29)$$

$$a_3 = m + r_1 \quad (30)$$

Equations (27) to (30) are then substituted into the solution of the general quartic of the previous section, which gives another twelve roots, x_{mn} . This time four of these roots satisfy both the quartic (2) with coefficients (27) to (30) and the quintic (1), let them be r_2, r_3, r_4 and r_5 . They are determined in the same way as r_1 , using nested “for loops” and “if-then” statements on Maple. The the final array has five non-zero elements, they are the five complex roots of the quintic equation,(1).

5 Conclusion

The root ambiguity is removed from the solution to the general fifth and fourth-degree polynomials.

6 Reference

1) Drociuk, Richard, “On the Complete Solution to the Most General Fifth Degree Polynomial”, GM/0005026v1 3 May 2000.